

Detection of *Chlamydophila psittaci* from pigeons by polymerase chain reaction in Ahvaz

Masoud Ghorbanpoor¹, Naghmeh-Moori Bakhtiari^{1*}, Mansoor Mayahi², Hana Moridveisi³

¹Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran. ²Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran. ³Graduated student of Veterinary Faculty, Shahid Chamran University of Ahvaz, Iran.

Received: July 2014, Accepted: January 2015

ABSTRACT

Background and Objective: *Chlamydophila psittaci* is a lethal bacterium that causes endemic avian chlamydiosis, and respiratory psittacosis. Laboratory diagnosis of *Chlamydophila psittaci* is difficult by culture. This study was designed to investigate the presence of *Chlamydophila psittaci* in collected pharyngeal swabs from asymptomatic pigeons by PCR.

Materials and Methods: Pharyngeal samples from pigeons with no symptoms of disease (n=280) were collected during hot and cold seasons in different parts of Ahvaz. DNA was extracted from specimens and subjected to PCR targeting *pmp* genes and 16s-23s rRNA intergenic spacer of *Cp. psittaci* and *Chlamydiales* specific primers.

Results: Of 280 samples 2 (0.7%) harbor were positive for *chlamydiales* (16s-23s intergenic spacer) and *Cp. psittaci* specific genes (*pmp* gene).

Conclusions: In this research the pigeons were asymptomatic carriers for *Cp. psittaci* in their respiratory discharges. These results suggest that *Cp. psittaci* infection of human can occur in very close and continuous contact with pigeons.

Keywords: *Chlamydophila psittaci*, Pigeon, PCR

INTRODUCTION

The *Chlamydiaceae* family is composed of a group of obligate intracellular bacteria that are found worldwide. This family is classified in two genera: the genus *Chlamydia* (including the species *C. muridarum*, *C. suis* and *C. trachomatis*) and the genus *Chlamydophila* (including the species *Cp. abortus*, *Cp. caviae*, *Cp. felis*, *Cp. pecorum*, *Cp. pneumoniae* and *Cp. psittaci*) (1). *Chlamydophila psittaci* (*Cp. psittaci*), can infect 465 avian species in

30 avian orders, with at least 153 species in the order Psittaciformes (2). This bacterium can be transmitted as metabolically inactive particles called elementary bodies (EBs) (3) from pet birds to humans (4). Transmission of this atypical respiratory pathogen can occur through direct contact with infected birds, birds' feces, the cloacae, nasal discharges, conjunctiva secretions causing respiratory disease in both mammals and birds (5). Transmission of *Cp. psittaci* from birds to humans, particularly to people at high risk like veterinarians, bird breeders and animal shopkeepers, has been reported before (6-8).

Chlamydophila psittaci has seven known genotypes (A-F and E/B) (9) and two non avian genotypes (M56 and WC). All genotypes can be transmitted to humans and cause psittacosis or parrot fever. Genotypes are distinguished by sequencing of the outer membrane protein A (*ompA*) gene (10). *Cp. psittaci* has the ability to remain infectious in the environment for months, presenting a variety of public health issues,

*Corresponding author: Naghmeh Moori Bakhtiari, DVM, PhD

Address: Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University, Golestan Street, Ahvaz, Iran.

Tel: +98-611-3333010-19

E-mail: n.moori@scu.ac.ir

including economically devastating outbreaks in poultry farms and occasionally severe pneumonia in humans (11). Chlamydiosis' symptoms in humans are variable, ranging from no clinical signs at all to severe systemic disease (12).

According to OIE (World Organization for Animal Health) recommendations, avian chlamydiosis diagnosis requires either isolation and identification of the organism, or demonstration of a four-fold increase in specific humoral antibody, as well as typical clinical signs (13). Amongst the tools available to demonstrate chlamydial presence in the host, PCR is an applicable technique due to its high specificity and sensitivity. This method presents the advantages of being simple, fast, easy to standardize and more suitable and safer than culture for processing large numbers of specimens (16).

The available current conventional PCR protocols for detection of avian species use single copy genes such as the *ompA* gene (2-11,14) or ribosomal RNA genes (16S-23S) (15) as amplification targets. More recently, a new set of primers (CpsiA/CpsiB) targeting conserved *pmp* (polymorphic membrane protein) genes family of *C. abortus* has been suggested as an effective and improved tool for PCR detection of avian chlamydiosis (16).

This family of genes, unique to *Chlamydiaceae*, was highlighted in all the genomes sequenced to date. Even though their function is still unknown, Pmp proteins are predicted to be localized in the outer membrane. In this study, the use of the CpsiA/CpsiB primers was extended to represent chlamydial strains of the six major serovars (serovars A-F) in the samples collected in field.

The aim of this study was to determine the presence of *Cp. psittaci* in pigeon population in Ahvaz city, Khuzestan province by amplification of *chlamydiales* and *Cp. psittaci* species specific genes in PCR.

MATERIALS AND METHODS

Sampling and DNA extraction. In the present study, 280 samples of pharyngeal swab from chana fossa of pigeons in transport medium (18) in different areas of Ahvaz were collected during two seasons (hot and cold) and analyzed by PCR for detection of *pmp* genes of *Cp. psittaci*. Genomic DNA was extracted from swab samples. Pharyngeal swab material in transport medium were homogenized in 0.5 ml TE buffer (10mM Tris-hydrochloric acid, 1mM EDTA,




Fig. 1. Results of the PCR assay for identification of *Cp. psittaci* gene

pmp:

- : negative control

+ : positive control strains of *Cp. psittaci*

Numbers 1,2: positive samples with *pmp* gene (300bp)

Numbers 3,4,5,6,7,8: negative samples without *pmp* gene

LA: 100bp ladder.




Fig. 2. Results of the PCR assay for identification of chlamydiales 16-23S spacer gene.

from right to left:

- Ladder 100bp

- Negative control

- Positive control strains of *Cp. psittaci* (352bp)

- Two positive samples with 16-23S spacer gene.

pH 8) supplemented with dithiothreitol (2 percent) and boiled for 10 minutes. After centrifugation, aliquots of 50 to 100 μ l of the supernatant were used in the PCR (19). The extracted DNA of each sample was kept frozen at -20°C until used. *Cp. psittaci* strain 6BC was used as positive control and a negative DNA control was performed by adding 1 μ l of sterile ultrapure deionized water.

Gene amplification. cIGS-1f/cGIS-2r (20) and CpsiA/CpsiB (2-16) protocols, which target 16S-23S rRNA intergenic spacer (*chlamydiales*) and *pmp* genes (*Cp. psittaci*), respectively, were used in this study. Genomic DNA purified from biological samples was firstly submitted to a CpsiA/CpsiB PCR

Table 1: Genes studied and specific primers in this research

Reference	Size (bp)	Sequence	Gene
Borel et al. 2006a	352	Forward: CAA GGT GAG GCT GAT GAC5'- Reverse: TCG CCT KTC AAT GCC AAG5'-	Chlamydiales (16-23S spacer)
Laroucau et al. 2001	300	Forward: ATG AAA CAT CCA GTC TAC TGG5'- Reverse: TTG TGT AGT AAT ATT ATC AAA5'-	Chlamydophila psittaci (gene pmp)
Binnicker et al. 2004	488	Forward :TCA AGA ACG AAA GTC GGA GG5'- Reverse :GGA CAT CTA AGG GCA TCA CA5'-	eukaryotic rRNA18 s

amplification including *Cp. psittaci* 6BC strain DNA in the mix (equivalent to 100 CFU per reaction) in order to assess the presence of PCR inhibitors in these samples. For checking the DNA extraction protocol 16S-23S rRNA intergenic spacer and 18s rRNA primers were used for prokaryotic and eukaryotic ribosomal RNA, respectively (Table1).

RESULTS

Using prokaryotic and eukaryotic ribosomal RNA specific primers, several DNA extracted from pharyngeal swabs and white blood cells were examined separately in PCR.

Out of 280 pharyngeal samples, 2 (0.71%) were positive for *Cp. psittaci* infection using PCR. Analysis of PCR products for presence of *pmp* gene of *Cp. psittaci* on agarose gel revealed a 300 bp fragment (Fig. 1). The positive control showed the expected amplification product specific for *Cp. psittaci* (300 bp). The chlamydiales specific primers (cIGS-1f/cGIS-2r) amplified a 352bp fragment in positive samples and the positive control (Fig. 2). Since positive results have been obtained by PCR from asymptomatic birds, it has been concluded that pharyngeal swab is suitable for detection of chlamydial infections. The relation between season, age, and gender with PCR results have been shown in Table 2.

In this study the pigeons were asymptomatic

carriers for *Cp. psittaci* in their respiratory excretions.

DISCUSSION

The *Chlamydiaceae* are etiological agents of many important human and animal diseases (21). *Chlamydophila psittaci* is a causative agent of psittacosis, systemic diseases in psittacine birds which can be acute, protracted, chronic, or subclinical manifestation that represents the most important animal chlamydiosis of zoonotic character (22). Furthermore, from 1941 to 2003, 78 cases in humans (23) and from 2005 to 2009, 66 human cases of psittacosis were reported to the Centers for Disease Control and Prevention (CDC) due to contact with feral pigeons (2-5). Polymerase chain reaction has been documented as a highly sensitive method for the detection of *Cp. psittaci* (25, 26). *CpsiA/CpsiB*, which were primarily designed from the *Cp. abortus* *pmp* genes (27) and described as being able to detect four different avian strains (2-16), are suitable for amplification of the 7 *pmp* related *Cp. psittaci* 6BC genes (data not shown). The total number of *pmp* genes for *Cp. psittaci* is still unknown. With a set of primers such as *CpsiA/CpsiB*, which targets several genomic fragments, the bacterial dilution effect is limited. Besides, the short size of the PCR products is an additional advantage for achieving better sensitivity results. All of the samples were amplified

Table 2. Analysis of results based on gender, age and season

PCR		Season	PCR		Age	PCR		Gender
Negative	Positive		Negative	Positive		Negative	Positive	
138	2	Hot	167	2	1>	156	1	Male
140	-	Cold	111	-	1<	122	1	Female
278	2	Total	278	2	Total	278	2	Total

with CpsiA/CpsiB primers. On clinical samples, designed primers are the best among those tested for detection of *Cp. psittaci* by simple conventional PCR. RFLP experiments performed using PCR fragments amplified with the CpsiA/CpsiB primers gave promising results, demonstrating that these primers may provide an interesting tool for molecular typing when the bacterium cannot be grown from pathological samples (2-17). In several studies, polymerase chain reaction and PCR-based methods were used to detect chlamydial DNA in pharyngeal, faecal, genital secretion samples and semen (28). A nested PCR-enzyme immunoassay (PCR-EIA) was developed to detect the *Cp. psittaci* outer membrane protein A (*ompA*) gene in pharyngeal swabs. The *ompA* nested PCR-EIA was more sensitive than the 16S-rRNA based nested PCR and isolation, revealing 105 out of 200 (52.5%) positives against 13 and 74 for the latter two tests, respectively. Teankum *et al.* examined the prevalence of chlamydial infection in semen and genital tracts of boars. They evaluated suitable PCR assays with bacteriology, LPS-ELISA, and immunohistochemistry (IHC). Three Chlamydia specific PCR assays targeting the 16S rRNA gene and the flanking IGS were compared. DNA dilution experiments revealed that 16Sig and IGS-S Chlamydia-specific PCRs were at least 10-fold more sensitive than the IGS-L PCR and serological and immunohistochemistry techniques (29).

Olsen *et al.* (1998) examined 312 fecal samples from 18 bird species to investigate to what extent wild passerine birds are carriers of *Cp. psittaci*. By using the PCR technique and subsequent DNA sequencing, they demonstrated *Cp. psittaci* DNA in fecal samples from nine (2.9%) birds of six different species. In this study, *Cp. psittaci* DNA was detected in 2 (1.4%) of 280 pharyngeal samples. The technique offers advantages over other testing methods in that it is fairly rapid and does not require organisms to be present in large numbers or to be viable (30).

Heddema *et al.* determined the prevalence and genotype of *Cp. psittaci* in fresh fecal samples from feral pigeons in Amsterdam, The Netherlands. The prevalence was 7.9% overall (26/331; 95% confidence interval, 5 to 11). Ten genotyped PCR-positive samples were all genotype B (31). Also, Sareyyupoglu *et al.* investigated the shedding of *Cp. psittaci* in fecal samples from 47 cage birds using PCR testing in Turkey. Following PCR with *Cp. psittaci* specific primers, 43 (91.5%) samples

were determined to harbour-specific DNA (32). Doosti and Arshi (2011) determined the prevalence of *Chlamydophila psittaci* in pigeon feces in Iran using PCR assay. DNA was extracted from 445 fecal samples of pigeons. The prevalence of this pathogen was 14.3% in the region of this study (33).

In a study conducted by the Pantchev *et al.* (2009), real-time PCR assays revealed 71 cases (92.2%) of *Cp. psittaci* infection in specimens from parrots and parakeets. Therefore, a suitable method was designed to research on chlamydial infections (34). The difference observed in the results can be related to host species (wild or domestic), physiological status (with or without symptoms), maintenance (free or cage), sampling, technique and geographical area.

In conclusion, we detected a lower prevalence of *Cp. psittaci* DNA from pharyngeal sample of pigeons with PCR assay in Ahvaz, which indicates the risk for veterinarians, employees and operators of aviaries and pet shops, and pet owners regarding exposure to the disease. This is the first study in Iran that investigated the shedding of *Cp. psittaci* in pharyngeal samples using PCR. We hope it will provide an alternative and easier method for the routine diagnostic laboratories working in the field of avian medicine.

ACKNOWLEDGEMENT

We thank Prof. Baris Sareyyupoglu (Department of Microbiology, Faculty of Veterinary Medicine, Ankara, Turkey) for his kind support in providing us the control *Cp. psittaci* DNA.

REFERENCES

1. Everett KD, Andersen AA. Identification of nine species of the Chlamydiaceae using PCR-RFLP. *Int J Syst Bacteriol* 1999; 49 (Pt 2); 803-813.
2. Vanrompay D, Harkinezhad T, van de Walle M, Beeckman D, van Droogenbroeck C, Verminnen K *et al.* *Chlamydophila psittaci* transmission from pet birds to humans. *Emerging Infectious Diseases* 2007; 13:1108-1110.
3. Binet R, Maurelli AT. Frequency of development and associated physiological cost of azithromycin resistance in *Chlamydia psittaci* 6BC and *C. trachomatis* L2. *Antimicrob Agents Chemother* 2007; 51: 4267-4275.
4. Johnston WB, Eidson M, Smith KA & Stobierski MG. Compendium of chlamydiosis (psittacosis) control, 1999. Psittacosis Compendium Committee, National Association of State Public Health Veterinarians. *J Am Vet Med Assoc* 1999; 214:640-646.

5. Smith, KA, Bradley KK, Stobierski MG, Tengelsen, LA. Compendium of measures to control *Chlamydomphila psittaci* (formerly *Chlamydia psittaci*) infection among humans (psittacosis) and pet birds. *J Am Vet Med Assoc* 2005; 226: 532-539.
6. Huminer D, Samra Z, Weisman Y, Pitlik S. Family outbreaks of psittacosis in Israel. *Lancet* 1988; 2 (8611):615-618.
7. Hinton DG, Shipley A, Galvin JW, Harkin JT, Brunton RA. Chlamydiosis in workers at a duck farm and processing plant. *Aust Vet J* 1993; 70: 174-176.
8. Saito T, Ohnishi J, Mori Y, Iinuma Y, Ichiyama S, Kohi F. Infection by *Chlamydomphila avium* in an elderly couple working in a pet shop. *J Clin Microbiol* 2005; 43: 3011-3013.
9. Geens T, Desplanques A, Van Loock M, Bonner BM, Kaleta EF, Magnino S. Sequencing of the *Chlamydomphila psittaci* ompA gene reveals a new genotype, E/B, and the need for a rapid discriminatory genotyping method. *J Clin Microbiol* 2005; 43: 2456- 2461.
10. Vanrompay D, Butaye P, Sayada C, Ducatelle R, Haesebrouck F. Characterization of avian *Chlamydia psittaci* strains using *omp1* restriction mapping and serovar-specific monoclonal antibodies. *Res Microbiol* 1997; 148:327-333.
11. Kaltenboeck B, Kousoulas KG, Storz J. Detection and strain differentiation of *Chlamydia psittaci* mediated by a twostep polymerase chain reaction. *J Clin Microbiol* 1991; 29 : 1969-1975.
12. Andersen AA, Vanrompay D. Avian chlamydiosis. *Rev Sci Tech* 2000; 19: 396-404.
13. OIE. Manual of diagnostic test and vaccines for terrestrial animals (mammals, birds and bees). 2004, Fifth edition.
14. Sayada C, Andersen AA, Storey CC, Milon A, Hashimoto N, Hirai K, et al. Usefulness of *omp1* restriction mapping for avian *Chlamydia psittaci* isolate differentiation. *Res Microbiol* 1995; 146: 155-165.
15. Everett KD, Andersen AA. The ribosomal intergenic spacer and domain I of the 23S rRNA gene are phylogenetic markers for *Chlamydia* spp. *Int J Syst Bacteriol* 1997; 47: 461-473.
16. Laroucau K, Souriau A and Rodolakis A. Improved sensitivity of PCR for *Chlamydomphila* using *pmp* genes. *Vet Microbiol* 2001; 82: 155-164.
17. Laroucau K, Trichereau A, Vorimore F, Mahe AM. A *pmp* genes-based PCR as a valuable tool for the diagnosis of avian chlamydiosis. *Vet Microbiol* 2006; 121: 150-157.
18. Quinn PJ, Carter ME, Markey B, Carter GR. *Clin Vet Microbiol* 2004, Appendix 2, pp:624.
19. Hewinson RG, Rankin SES, Bevan BJ, Field M, Woodward MJ. *Vet Rec* 1991; 128: 129-130.
20. Borel N, Mukhopadhyay S, Kaiser C, Sullivan ED, Miller RD, Timms P, et al. Tissue Micro-Array (TMA) analysis of normal and persistent *Chlamydomphila pneumonia* infection. *BMC Infect Dis* 2006a; 6: 152.
21. Bush RM, Everett KD. Molecular evolution of the Chlamydiaceae. *Int J Syst Evol Microbiol* 2001: 51: 203-220.
22. Goellner S, Schubert E, Liebler-Tenorio E, Hotzel H, Saluz HP, Sachse K. Transcriptional response patterns of *Chlamydomphila psittaci* in different in vitro models of persistent infection. *Infect Immun* 2006; 74: 4801-4808.
23. Haag-Wackernagel D, Moch H. Health hazards posed by feral pigeons. *J Infect* 2004; 48: 307-313.
24. Huminer D, Pitlik S, Kitayin D, Weissman Y, Samra Z. Prevalence of *Chlamydia psittaci* infection among persons who work with birds. *Isr J Med Sci* 1992; 28:739-741.
25. Takashima I, Imai Y, Itoh N, Kariwa H, Hashimoto N. Polymerase chain reaction for the detection of *Chlamydia psittaci* in the feces of budgerigars. *Microbiol Immunol* 1996; 40: 21-26.
26. Trevejo RT, Chomel BB and Kass PH. Evaluation of the polymerase chain reaction in comparison with other diagnostic methods for the detection of *Chlamydia psittaci*. *J Vet Diagn Invest* 1999; 11: 491-496.
27. Longbottom D, Russell M, Jones G, Lainson F, Herring A. Identification of a multigene family coding for the 90 kDa proteins of the ovine abortion subtype of *Chlamydia psittaci*. *FEMS Microbiol Lett* 1996;142: 277-281.
28. Van Loock M, Verminnen K, Messmer TO, Volckaert, G, Goddeeris BM, and Vanrompay D . Use of a nested PCR-enzyme immunoassay with an internal control to detect *Chlamydomphila psittaci* in turkeys. *BMC Infectious Diseases* 2005; 5:76.
29. Teankum K, Pospischil A, Janett F, Bu'rgi E, Brugnera E, Hoelzle K, Polkinghorne A, Weilenmann R, Zimmermann DR, Borel N. Detection of chlamydiae in boar semen and genital tracts. *Vet Microbiol* 2006; 116: 149-157.
30. Olsen B, Persson K and Broholm KA. PCR detection of *Chlamydia psittaci* in faecal samples from passerine birds in Sweden monoclonal antibodies. *Res Microbiol* 1998; 148: 327-333.
31. Heddema ER, Beld MGHM, de Wever B, Langerak AAJ, Pannekoek Y and Duim B. Development of an internally controlled real-time PCR assay for detection of *Chlamydomphila psittaci* in the LightCycler 2.0 system. *Clin Microbiol Infect* 2006; 12: 571-575.
32. Sareyyupoglu B, Cantekin Z and Bas B. *Chlamydomphila psittaci* DNA detection in the faeces of cage birds. *Zoonoses Public Health* 2007; 54: 237-242.
33. Doosti A and Arshi A. Determination of the Prevalence of *Chlamydia psittaci* by PCR in Iranian Pigeons. *Int J Biol* 2011; 3, 4: 79-82.
34. Pantchev P, Sting R, Bauerfeind R, Tyczka J, Sachse k. new real-time PCR tests for species-specific detection of *chlamydomphila psittaci* and *chlamydomphila abortus* from tissue samples. *Vet J* 2009; 181: 145-150.